Imperial/TP/97-98/060
hep-th/9807097

arXiv:hep-th/9807097v4 6 Aug 1998

On Four-point Functions in the CFT/AdS Correspondence

Hong Liu* and A.A. Tseytlin*†

Theoretical Physics Group, Blackett Laboratory,
Imperial College, London SW7 2BZ, U.K.

Abstract

We discuss the properties of four-point functions in the context of the correspondence between a classical supergravity theory in the bulk of the Anti de Sitter (AdS) space and quantum conformal field theory (CFT) at the boundary. The contribution to a four-point function from the exchange of a scalar field of arbitrary mass in AdS space is explicitly identified with that of the corresponding operator in the conformal partial wave expansion of a four-point function on the CFT side. Integral representations are found for the massless vector and graviton exchanges. We also discuss some aspects of the four-point functions of trF^2 and $trFF^*$ ('dilaton' and 'axion') operators in the $\mathcal{N}=4$ supersymmetric $SU(N)$ Yang-Mills theory as predicted by type IIB supergravity in the AdS_5 background.

July 1998

* e-mail address: hong.liu@ic.ac.uk

* e-mail address: tseytlin@ic.ac.uk

† Also at Lebedev Physics Institute, Moscow.

1. Introduction

There is a recent revival of interest in the connections between large N Yang-Mills theory [1] and string theory [2] following, in particular, the conjecture [3] that there is an exact correspondence between string/M theory on the Anti de Sitter space AdS_{d+1} and certain superconformal field theory (CFT_d) defined at the boundary of the AdS_{d+1} (see also [4]). According to the conjecture, the quantum $\mathcal{N} = 4$ supersymmetric Yang-Mills theory with gauge group $SU(N)$ in the large N and large 't Hooft coupling limit can be described by the classical type IIB supergravity on $AdS_5 \times S^5$ space.

The formulation of the conjecture was made more explicit in [5,6], where it was proposed that the partition function of supergravity/string theory with fixed boundary values of the fields is to be identified with the generating functional of the composite operators in the CFT. There is a one-to-one correspondence between certain local operators Φ_i of the boundary CFT and the bulk fields ϕ_i in AdS [5,7,6,8]. The boundary CFT operator Φ_i and the associated bulk field ϕ_i carry the same unitary, irreducible and the highest weight representation of the conformal group $SO(d, 2)$, where the scale dimension λ_i of Φ_i is identified with the lowest energy value of ϕ_i and can be further related to the mass of ϕ_i. The correlation functions of the CFT operators are identified with the classical "S-matrix elements" of the bulk fields with their boundary values fixed. Two-point and three-point functions follow simply from the quadratic terms and cubic vertices of the bulk theory, while the four-point functions, in general, contain the contact contributions as well as the exchanges of virtual particles.

Using this proposal, some 'model' and 'realistic' two-point and three-point functions have been computed in [9,10,11,12,13,14,15,16]. In particular, a family of three-point functions of chiral primary operators in $\mathcal{N} = 4$ SYM have been evaluated and shown to be equal to their free-field values, suggesting a non-renormalization theorem in large N limit [17].

In this paper we investigate the properties of the 4-point functions in the context of the CFT/AdS correspondence. Four-point functions from contact interactions (quartic vertices) were considered before in [10]. The scalar exchange diagrams with some special values of mass were also discussed in [18]. The exchange diagrams in AdS are, in gen-

eral, very difficult to evaluate explicitly, as the propagators and the integrals are quite complicated. Here we follow a different approach.

In CFT$_d$, the states generated by acting by a product of the conformal operators on the vacuum can be decomposed into a direct sum of irreducible representations of the conformal group

$$\Phi_1(x_1)\Phi_2(x_2)|0>=\sum_k \int d^d x\ Q_k(x|x_1,x_2)|k,x>\ , \tag{1.1}$$

where k sums over all the irreducible representations in the Hilbert space and states $|k,x>=\Phi_k(x)|0>$ span the space of irreducible representation T_k. This conformal partial wave expansion (CPWE) was obtained in early seventies by several authors [19,20] (see [21,22,23] for reviews). Using (1.1), the four-point functions can be written as

$$<0|\Phi_1(x_1)\Phi_2(x_2)\Phi_3(x_3)\Phi_4(x_4)|0>=\sum_k G_k\ , \tag{1.2}$$

where G_k is the contribution to the four-point function from the intermediate states generated by the operator Φ_k,

$$G_k=\int d^d x d^d y\, Q_k^*(x_1,x_2|x)<k,x|k,y>Q_k(y|x_3,x_4)\ . \tag{1.3}$$

Since the bulk propagator of the field ϕ_k in AdS_{d+1} associated to the conformal operator Φ_k by the AdS/CFT correspondence can be also written as a sum over normal modes in AdS_{d+1} which span the same irreducible representation T_k of the isometry/conformal group $SO(d,2)$, one is tempted to conjecture that G_k represents the contribution of the ϕ_k exchange diagram in AdS_{d+1}. Diagrammatically, this equivalence can be expressed as



Figure 1: Equivalence between CPWE and scattering in AdS.

One of the aims of the present paper is to prove that this is indeed the case for the intermediate states corresponding to the scalar operators. It should be possible to generalise our method also to operators of higher spin. We shall attempt to consider the two cases which are of particular interest: massless vector and massless tensor (graviton) exchanges. They correspond to the conserved current vector and the energy momentum tensor operators in the CFT. The explicit demonstration of the equivalence between the CPWE representation for the CFT correlator and the AdS amplitude here appears to be more difficult and will not be given in the present paper. The expressions for the propagators for the photon and graviton in AdS are quite involved (useful expressions for them which are suitable for explicit calculations were not previously given in the literature, cf. [24,25]). There are also complications related to the presence of the gauge degrees of freedom and the fact that the current and the energy momentum tensor carry indecomposable representations of the conformal group. In this paper we shall use a non-covariant gauge fixing and will be able to write down the AdS amplitudes with the photon and graviton exchanges in the complete integral form. The detailed analysis of these amplitudes and establishing their relation to CPWE will not be attempted here.

Having identified the exchange diagrams with CPWE, a question that naturally arises is the interpretation of contact interactions (quartic or higher vertices) on the CFT side, as there is no obvious counterpart for them in CPWE. One possibility is that since contact terms can always be formally written as special exchange diagrams, e.g.,

$$\int d^{d+1}x\, \phi_1\phi_2\phi_3\phi_4 = -\int d^{d+1}x\; d^{d+1}y\;\; \phi_1\phi_2\; G(x,y)\; \partial^2(\phi_3\phi_4)\;, \tag{1.4}$$

where G is the massless field propagator, $-\partial^2 G = \delta(x-y)$, they might be contained in CPWE. Another possibility could be that the boundary CFT is not closed. Let us consider for example the $\mathcal{N}=4$ SYM theory in the large N and large 't Hooft coupling $\lambda = g^2N$ limit. Suppose that a four-point function at finite λ is expanded in the form of (1.2) and (1.3). As we increase λ, certain correlators may approach zero as some inverse powers of λ and thus may not contribute in the $\lambda\to\infty$ limit. But if there is a very large number of such vertices, their total contribution to the sum (1.2) may not vanish. This would correspond

to the presence of contact interactions in supergravity.[1] As there are many contact terms in IIB supergravity on $AdS_5 \times S^5$, such possibility deserves a detailed investigation.

The existence of the AdS/CFT correspondence puts by itself strong constraints on the theories on both sides. Since conformal field theories always contain the energy-momentum tensor which generates the conformal algebra, the theory in AdS must be a gravitational theory. If the theory in AdS is a field theory (supergravity), then on CFT side the possible three-point functions and intermediate states contributing to (1.2) are highly constrained as only vertices which can be written as local invariants in AdS are allowed. For example, consider a correlation function of four scalars. In general, symmetric tensor operators of spin greater than two can contribute to it as intermediate states in (1.2). However, there is no local covariant interaction vertex for two scalars and a higher spin tensor in supergravity theory (though it may be present in string theory), so it should vanish also on the CFT side. Assuming the equivalence between CPWE in CFT and the scattering amplitude in AdS, we see also that different channels for CPWE in CFT should correspond to $s-t-u$ channels in the scattering amplitudes in AdS, which seems to imply that the scattering amplitudes in AdS should have $s-t-u$ crossing symmetry.[2]

In case of the "$\mathcal{N}=4$ SYM – IIB supergravity on $AdS_5 \times S^5$" correspondence the supergravity four-point functions in general are quite complicated.[3] In this paper we shall focus on the dilaton-axion sector, where the corresponding four-point functions are given by a relatively small number of diagrams. We will show that the main non-trivial

[1] These contact supergravity vertices may be thought of as originating from string field theory (with only cubic interactions between massless and massive modes) in the low-energy approximation in which all massive string modes are integrated out.

[2] One could think that this might be an indication that the theory on the AdS side should actually be a string-type theory. One does not expect to find the crossing symmetry in the bulk supergravity amplitudes but this is less clear when the bulk-to-boundary propagators are attached. Duality is, of course, restored in the bulk amplitudes once one replaces the supergravity amplitudes by the full string amplitudes, i.e. includes all α'-corrections. At the same time, the boundary theory at large N and large g^2N corresponding just to supergravity with no α'-corrections is also a CFT which should have CPWE.

[3] Higher-order α' and non-perturbative string-theory contributions to four-point functions in $\mathcal{N}=4$ large N SYM were discussed in [26].

contribution in this sector comes from a graviton exchange (the 'mixed' scalar four-point function contains also a contact contribution). Using the graviton propagator found here in the non-covariant $h_{0\mu} = 0$ gauge, we will be able to present the complete expressions for the scalar four-point functions in terms of the formal integrals. We shall discuss briefly certain properties of the integrals, leaving their evaluation and establishing the correspondence with the CPWE in CFT for future.

The structure of the paper is as follows. In section 2 we shall review some aspects of CFT in d dimensions, in particular, the conformal partial wave expansion for the four-point functions. In section 3 we shall discuss the scattering diagrams involving scalar exchanges in AdS space. We will show that they can be identified with the contributions to CPWE coming from the corresponding operators on the CFT side. In section 4 we shall study the scattering amplitudes involving exchanges of massless vector and graviton. In section 5 we shall consider the dilaton and axion four-point functions in $D = 5$ supergravity corresponding to the correlators of the trF^2 and $trFF^*$ operators in $\mathcal{N} = 4$ SYM theory. Appendix A contains notation and some technical details about the scalar propagator in AdS_{d+1}. In Appendix B we recall the expression [19] (see also [27]) for the scalar operator contribution to the scalar four-point function in CFT_d.

2. Four-point functions and conformal partial wave expansion in CFT

Let us first review certain aspects of the conformal field theory in d dimension [21,22,23]. Denote the space of an irreducible representation T_σ of the conformal group[4] as M_σ, $\sigma = (\lambda, \vec{s})$, where λ is the conformal dimension and $\vec{s}$ is a set of quantum numbers labelling the spin degrees of freedom. We assume that the Hilbert space can be represented as a direct sum of spaces M_{σ_i}, i.e.

$$\mathcal{H} = M_{\sigma_1} + M_{\sigma_2} + ... + M_{\sigma_i} + ... , \qquad \sigma_i = (\lambda_i, \vec{s}_i), \quad i = 1, 2, ... \tag{2.1}$$

[4] For the Minkowskian signature, the representations under consideration are those of the infinite covering group of $SO(d, 2)$ which are unitary and satisfy the spectrality condition: $p_0 > 0$, $p^2 > 0$. For the Euclidean signature, they are the irreducible representations of $SO(d + 1, 1)$ which can be analytically continued from the Minkowskian counterparts.

Conformal fields $\Phi_{\sigma_i}(x)$ are defined as the operators which generate spaces M_{σ_i},

$$M_{\sigma_i} = \{|\sigma_i, x> \text{ for all } x\} = \{\Phi_{\sigma_i}(x)|0> \text{ for all } x\}.$$

States of the type $\Phi_{\sigma_i}(x_1)\Phi_{\sigma_j}(x_2)|0>$ can be decomposed in terms of the basis in (2.1),

$$\Phi_{\sigma_i}(x_1)\Phi_{\sigma_j}(x_2)|0> = \sum_k \int d^d x \; Q_{ijk}(x|x_1, x_2)|\sigma_k, x> \quad . \tag{2.2}$$

This implies the operator product expansion (OPE)

$$\Phi_{\sigma_i}(x_1)\Phi_{\sigma_j}(x_2) = \sum_k \int d^d x \; Q_{ijk}(x|x_1, x_2)\Phi_{\sigma_k}(x) \; . \tag{2.3}$$

The standard OPE in the nearby points can be obtained from (2.3) by expanding the integrand in $y = x_1 - x_2$,

$$\Phi_{\sigma_i}(y)\Phi_{\sigma_j}(0)|_{y\to 0} \; \sim \; \sum_{k,m} A_{\sigma_k,m}(y)\Phi_{\sigma_k,m}(0) \; , \tag{2.4}$$

where $A_{\sigma_k,m} \sim y^{-(\lambda_i+\lambda_j-\lambda_k-m)}$ and $\Phi_{\sigma_k,m}$ are m-th order derivatives of the field Φ_{σ_k}.

When Φ_σ's are orthogonal to each other, Q's are just the amputated three-point functions,

$$Q_{ijk}(x|x_1, x_2) = \int d^d x' \; W^{-1}_{\sigma_k}(x - x') < 0|\Phi_{\sigma_k}(x')\Phi_{\sigma_i}(x_1)\Phi_{\sigma_j}(x_2)|0> \; , \tag{2.5}$$

with $W_\sigma(x - x') =< 0|\Phi_\sigma(x)\Phi_\sigma(x')|0>$ and its inverse W^{-1} defined by,

$$\int d^d x \, W_\sigma(x_1 - x)W^{-1}_\sigma(x - x_2) = I_+(x_1 - x_2) \; , \tag{2.6}$$

$$I_+(x) = \frac{1}{(2\pi)^d} \int d^d p \; \vartheta(p^0) \; \vartheta(p^2) \; e^{ipx} \; . \tag{2.7}$$

That I_+ (instead of the Dirac δ-function) appears on the right side of (2.6) follows from the spectrality condition.

States involving higher-order products of Φ_σ's can be written in the basis (2.1) by repeatedly using (2.3). The problem of solving the theory thus becomes equivalent to finding the spectrum and the couplings for the infinite set of fields Φ_{σ_i}.

Applying (2.3) to the four-point functions we find

$$W_{ijkl}(x_1,x_2,x_3,x_4) = <0|\Phi_{\sigma_i}(x_1)\Phi_{\sigma_j}(x_2)\Phi_{\sigma_k}(x_3)\Phi_{\sigma_l}(x_4)|0>$$
$$= \sum_m \int d^dx d^dy\ Q_{ijm}(x_1,x_2|x)\ W_{\sigma_m}(x-y)\ Q_{mkl}(y|x_3,x_4)\ . \tag{2.8}$$

We now switch to the Euclidean signature. The conformal partial wave expansion of the four-point function in Euclidean region takes the form[5] ($\sigma = (\lambda, \vec{s})$)

$$G(x_1,x_2,x_3,x_4) = \sum_{\vec{s}} \int_C d\lambda\ \hat{G}_\sigma(x_1,x_2,x_3,x_4)\ , \tag{2.9}$$

where the integral is a contour integral in the complex plane of the conformal dimension λ (see below) and

$$\hat{G}_\sigma(x_1,x_2,x_3,x_4) = n(\sigma)\int d^dx\ G_{ij\tilde{\sigma}}(x_1,x_2,x)G_{kl\sigma}(x,x_3,x_4)\ . \tag{2.10}$$

Here

$$G_{kl\sigma}(x,x_3,x_4) =< \Phi_\sigma(x)\Phi_k(x_3)\Phi_l(x_4) >\ , \qquad G_{ij\tilde{\sigma}}(x_1,x_2,x) =< \Phi_i(x_1)\Phi_j(x_2)\Phi_{\tilde{\sigma}}(x) >$$

are the Euclidean three-point functions with $\tilde{\sigma} = (d-\lambda, \vec{s})$ and $n(\sigma)$ is some normalization constant. The field $\Phi_{\tilde{\sigma}}$ is called the conformal partner (or shadow operator) of Φ_σ, related to it by[6]

$$\Phi_{\tilde{\sigma}}(x) = \int d^dy\ \Delta_{\tilde{\sigma}}(x-y)\Phi_\sigma(y)\ , \qquad \Delta_{\tilde{\sigma}}(x-y) =< \Phi_{\tilde{\sigma}}(x)\Phi_{\tilde{\sigma}}(y) >\ . \tag{2.11}$$

$\hat{G}_\sigma$ includes the contributions from both Φ_σ and its conformal partner $\Phi_{\tilde{\sigma}}$. The integration contour in (2.9) is chosen so that to select only the contribution from Φ_σ. One can decompose $\hat{G}_\sigma$ into the parts coming from Φ_σ and $\Phi_{\tilde{\sigma}}$,

$$\hat{G}_\sigma = G_\sigma + G_{\tilde{\sigma}}\ . \tag{2.12}$$

[5] We denote the Euclidean correlators by G and omit some subscripts to simplify the notation.

[6] Note that there is no spectrality condition in Euclidean space so that $\Delta_{\tilde{\sigma}}$ and Δ_σ can be chosen to satisfy

$$\int d^dx\ \Delta_\sigma(x_1-x)\Delta^{-1}_{\tilde{\sigma}}(x-x_2) = \delta(x_1-x_2)\ .$$

Since G_σ and $G_{\tilde{\sigma}}$ have different pole structure ($x_1 - x_2 \to 0$)

$$G_{ij\sigma} \sim \frac{1}{|x_1 - x_2|^{\lambda_i+\lambda_j-\lambda}} , \qquad G_{ij\tilde{\sigma}} \sim \frac{1}{|x_1 - x_2|^{\lambda_i+\lambda_j+\lambda-d}} ,$$

the decomposition (2.12) is unique. Using (2.12) instead of the contour integral we can rewrite (2.9) as

$$G(x_1, x_2, x_3, x_4) = \sum_{\sigma_i} G_{\sigma_i}(x_1, x_2, x_3, x_4) . \tag{2.13}$$

Besides (2.8) and (2.9), there are two other ways to write partial wave expansions: in terms of "u-channel" and "t-channel". The equivalence of the three channels is guaranteed by the associativity of the operator algebra (2.3). This is usually called the *crossing symmetry* of the four-point functions. As was already mentioned in the Introduction, the crossing symmetry of the CFT four-point functions should have interesting implications for the structure of the corresponding scattering amplitudes in the AdS space.

3. Four-point functions in CFT/AdS correspondence: scalar exchange

In this section we consider the contribution to a four-point function of the exchange of a scalar field of an arbitrary mass. For definiteness, we shall study the 'model' four-point functions of scalar fields and consider scattering in AdS resulting from vertices of the type $\phi\phi_1\phi_2$ and $\phi\phi_3\phi_4$. The scattering amplitude is given by (see Appendix A for our notations)

$$S_\nu(x_1, x_2, x_3, x_4) = \int \frac{du_0 d^d u}{u_0^{d+1}} \frac{dv_0 d^d v}{v_0^{d+1}} \mathcal{K}_{\lambda_1}(u, x_1)\mathcal{K}_{\lambda_2}(u, x_2)G(u, v)\mathcal{K}_{\lambda_3}(v, x_3)\mathcal{K}_{\lambda_4}(v, x_4) , \tag{3.1}$$

where $\mathcal{K}_\lambda(u, x)$ is the boundary propagator [6] corresponding to a conformal field with dimension λ

$$\mathcal{K}_\lambda(u, x) = c_\lambda(\frac{u_0}{|u - x|^2})^\lambda , \tag{3.2}$$

and $G(x, y)$ is the bulk propagator for a scalar field of mass m (see Appendix A) [10],

$$G(x, y) = (x_0 y_0)^{d/2} \int \frac{d^d k}{(2\pi)^d} e^{i\vec{k}\cdot(\vec{x}-\vec{y})} I_\nu(kx_0^<)K_\nu(kx_0^>) . \tag{3.3}$$

Here I and K are the modified Bessel functions, the parameter ν is related to the mass by $\nu = \sqrt{m^2 + \frac{1}{4}d^2}$, and $x_0^<(x_0^>)$ is the smaller (larger) number among x_0 and y_0. The

amplitude (3.1) with (3.3) inserted is not manifestly conformally invariant. Its conformal invariance can be seen by using an alternative representation for $G(x,y)$ [24],

$$G(x,y) = rz^{-\lambda} F(\lambda, \nu - \frac{1}{2}; 2\nu + 1, z^{-1}) \ , \tag{3.4}$$

where r is a normalization constant, F is a hypergeometric function and $z = \frac{(x_0+y_0)^2+(\vec{x}-\vec{y})^2}{4x_0 y_0}$. It is clear from (3.4) that $G(x,y)$ is invariant under transformation $x \to \frac{x}{|x|^2}$, $y \to \frac{y}{|y|^2}$.

Let us first look at the "pseudo-propagator", given by

$$\hat{G}(x,y) = (x_0 y_0)^{d/2} \int \frac{d^d k}{(2\pi)^d} e^{i\vec{k}\cdot(\vec{x}-\vec{y})} K_\nu(kx_0) K_\nu(ky_0) \ . \tag{3.5}$$

The value of (3.1) with (3.5) inserted instead of (3.3) will be denoted $\hat{S}$. After the substitution of boundary propagators and (3.5) it can be written as

$$\hat{S_\nu}(x_1,x_2,x_3,x_4) = \int \frac{d^d k}{(2\pi)^d} F^*_{12}(k;\vec{x}_1,\vec{x}_2) F_{34}(k;\vec{x}_3,\vec{x}_4) \ , \tag{3.6}$$

where

$$F_{12}(k;\vec{x}_1,\vec{x}_2) = \int \frac{du_0 d^d u}{u_0^{d+1}} u_0^{\frac{d}{2}} e^{-i\vec{k}\cdot\vec{u}} \ K_\nu(ku_0)\, \mathcal{K}_{\lambda_1}(u,x_1) \mathcal{K}_{\lambda_2}(u,x_2) \ . \tag{3.7}$$

The Fourier transformation for the boundary propagator with $\lambda = \nu + \frac{d}{2}$ is

$$\mathcal{K}_\lambda(u,x) = c_\lambda (\frac{u_0}{|u-x|^2})^\lambda = \int \frac{d^d k}{(2\pi)^d} \ e^{i\vec{k}\cdot(\vec{u}-\vec{x})} f(k,u_0) \ , \tag{3.8}$$

with

$$f(k,u_0) = \frac{1}{b_\lambda} u_0^{\frac{d}{2}} k^\nu K_\nu(ku_0) \ , \qquad b_\lambda = 2^{\nu-1}\Gamma(\nu) \ .$$

Thus

$$u_0^{\frac{d}{2}} K_\nu(ku_0) = b_\lambda k^{-\nu} \int d^d x \ e^{-i\vec{k}\cdot(\vec{x}-\vec{u})} \mathcal{K}_\lambda(u,x) \ .$$

Plugging this into (3.7), we find

$$F_{12}(k;\vec{x}_1,\vec{x}_2) = b_\lambda k^{-\nu} \int \frac{du_0 d^d u}{u_0^{d+1}} \int d^d x \ e^{-i\vec{k}\cdot\vec{x}} \mathcal{K}_\lambda(u,x) \mathcal{K}_{\lambda_1}(u,x_1) \mathcal{K}_{\lambda_2}(u,x_2)$$

$$= b_\lambda k^{-\nu} \int d^d x \ e^{-i\vec{k}\cdot\vec{x}} G_{\lambda\lambda_1\lambda_2}(x,x_1,x_2) = b_\lambda k^{-\nu} G_{\lambda\lambda_1\lambda_2}(k,x_1,x_2) \ ,$$

where

$$G_{\lambda\lambda_1\lambda_2}(x,x_1,x_2)=\int\frac{du_0d^du}{u_0^{d+1}}\mathcal{K}_\lambda(u,x)\mathcal{K}_{\lambda_1}(u,x_1)\mathcal{K}_{\lambda_2}(u,x_2) \tag{3.9}$$

is the three-point function according to the CFT/AdS correspondence. Similarly, we find

$$F_{34}(k;\vec{x}_3,\vec{x}_4)=b_\lambda k^{-\nu}G_{\lambda\lambda_3\lambda_4}(k,x_3,x_4)\ .$$

Then it follows from (3.6) that

$$\hat{S}_\nu=b_\lambda^2\int\frac{d^dk}{(2\pi)^d}\,G^*_{\lambda\lambda_1\lambda_2}(k,x_1,x_2)k^{-2\nu}G_{\lambda\lambda_3\lambda_4}(k,x_3,x_4)\ . \tag{3.10}$$

From (3.8) and (3.9) we can see that

$$G_{\tilde{\lambda}\lambda_1\lambda_2}(k,x_1,x_2)=\frac{b_\lambda}{b_{\tilde{\lambda}}}k^{-2\nu}G_{\lambda\lambda_1\lambda_2}(k,x_1,x_2) \tag{3.11}$$

Thus (3.10) can be written as

$$\hat{S}_\nu=b_\lambda b_{\tilde{\lambda}}\int\frac{d^dk}{(2\pi)^d}\,G^*_{\tilde{\lambda}\lambda_1\lambda_2}(k,x_1,x_2)G_{\lambda\lambda_3\lambda_4}(k,x_3,x_4)\ ,$$

and in coordinate space it becomes,

$$\hat{S}_\nu=b_\lambda b_{\tilde{\lambda}}\int d^dx\ \ G_{\lambda_1\lambda_2\tilde{\lambda}}(x_1,x_2,x)G_{\lambda\lambda_3\lambda_4}(x,x_3,x_4)\ . \tag{3.12}$$

We notice that the above expression (3.12) for $\hat{S}_\nu$ is precisely the same as the CFT expression (2.10) with $\sigma=(\lambda,0)$. Thus we have identified the amplitude $\hat{S}_\nu$ with the CFT correlator $\hat{G}_\sigma$ in (2.9).

Let us now look at the relation between S_ν (3.1) and $\hat{S}_\nu$ (3.6). Using that

$$K_\nu=\frac{\pi}{2}\frac{1}{\sin\nu\pi}(I_{-\nu}-I_\nu)\ ,$$

and

$$\int_0^\infty dx_0\int_0^\infty dy_0=\int_0^\infty dx_0\int_0^{x_0}dy_0+\int_0^\infty dy_0\int_0^{y_0}dx_0\ ,$$

it is easy to see that

$$\hat{S}_\nu=S_\nu+S_{-\nu}\ , \tag{3.13}$$

which can be understood as the sum of the contributions from the fields of dimensions λ and $d-\lambda$ respectively. Comparing (3.13) with (2.12) and assuming that S_ν has analytic dependence on ν, we find that with $\hat{S}_\nu$ identified with $\hat{G}_\sigma$ in (2.10), S_ν and $S_{-\nu}$ are equal to G_σ and $G_{\tilde{\sigma}}$, demonstrating the required relation between the AdS amplitude S_ν and the CFT correlator.

It is easy to see that the above procedure applies without change to other types of 3-point interaction (e.g., $\phi\partial\phi_1\partial\phi_2$) and to scattering of higher spin fields involving scalar exchange. Thus we have established the correspondence between the exchange diagrams in AdS and the conformal partial wave expansion in the CFT for the case of the *scalar* intermediate states.

We mention here that the contribution of a scalar operator to the CPWE (eqs (2.8), (2.9)) of a four-point function has been evaluated long time ago in [19]. The expression can be written in a closed form in terms of double hypergeometric functions (see Appendix B). From the identification of (3.1) and (2.8), we see that it can also be interpreted as the scattering amplitude (3.1) in AdS.

4. Photon and graviton propagators in AdS_{d+1}

In this section we shall consider the scattering amplitudes in AdS involving massless vector and graviton exchanges. The photon and graviton propagators in covariant gauges were discussed before in [25]. The expressions found were complicated and do not seem to be useful in explicit calculations. Here we shall choose the non-covariant gauges, $A_0 = 0$ for the vector and $h_{0\mu} = 0$ for the graviton. It turns out that the resulting AdS propagators are quite simple and have structure similar to that of their flat space counterparts.

4.1. Massless vector

Let us start with the case of a vector field in AdS space described by the action

$$I = \int d\alpha \, (\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + A_\mu \mathcal{J}^\mu) \ , \qquad \int d\alpha = \int d^{d+1}x \sqrt{g_0} \ . \tag{4.1}$$

We fix the Coulomb gauge $A_0 = 0$. In this gauge the equation for A_i $(i = 1, ..., d)$ becomes

$$(\partial_0^2 + \partial_j^2) A_i - \partial_j \partial_i A_j - \frac{d-3}{x_0} \partial_0 A_i = -\frac{1}{x_0^2} \mathcal{J}_i \ . \tag{4.2}$$

The equation for A_0 gives the constraint

$$\partial_0 \partial_i A_i = \frac{1}{x_0^2} \mathcal{J}_0 \ . \tag{4.3}$$

We decompose A_i as

$$A_i = A_i^\perp + \partial_i \xi \ ,$$

where $A_i^\perp$ is the transverse part and $\xi = \frac{1}{\partial^2} \partial_i A_i$ with $\partial^2 = \partial_i \partial_i$. Then (4.2) and (4.3) reduce to

$$\partial_0 \xi = \frac{1}{x_0^2} \frac{1}{\partial^2} \mathcal{J}_0, \qquad (\partial_0^2 + \partial^2) A_i^\perp - \frac{d-3}{x_0} \partial_0 A_i^\perp = -\frac{1}{x_0^2} \mathcal{J}_i^\perp \ ,$$

where $\mathcal{J}_i^\perp$ is the transverse part of $\mathcal{J}_i$, i.e. $\partial_i \mathcal{J}_i^\perp = 0$. Thus the equation for $x_0 A_i^\perp$ reduces to that for a free scalar of $m^2 = -(d-1)$ in AdS_{d+1} space with a source $x_0 \mathcal{J}_i^\perp$. It can be easily solved to obtain

$$x_0 A_i^\perp(x) = \int d\alpha \ \ G(x,y) \ y_0 \mathcal{J}_i^\perp(y) \ , \tag{4.4}$$

where $G(x,y)$ the propagator for a free scalar with $m^2 = -(d-1)$.

Then

$$I = \frac{1}{2} \int d\alpha \, A_i \mathcal{J}^i = \frac{1}{2} \int d\alpha \, (A_i^\perp \mathcal{J}^i + \partial_i \xi \mathcal{J}^i) \ . \tag{4.5}$$

Note that the conservation of the current gives

$$D_\mu \mathcal{J}^\mu = 0 \implies \partial_i \mathcal{J}^i = -\frac{1}{\sqrt{g_0}} \partial_0 (\sqrt{g_0} \mathcal{J}^0) \ .$$

After a partial integration, use of the current conservation and the substitution of (4.4) into (4.5) we get,

$$I = \frac{1}{2} \int \frac{dx_0 d\vec{x}}{x_0^{d+1}} \frac{dy_0 d\vec{y}}{y_0^{d+1}} \ x_0 y_0 \ \mathcal{J}_i^\perp(x) \ G(x,y) \ \mathcal{J}_i^\perp(y) - \frac{1}{2} \int \frac{dx_0 d\vec{x}}{x_0^{d+1}} \ \mathcal{J}_0 \frac{1}{\partial^2} \mathcal{J}_0 \ . \tag{4.6}$$

This expression determines the photon propagator.[7]

[7] In (4.6) there is also a boundary term $-\frac{1}{2} \int_{\partial M} d^d x \ x_0^{1-d} \ \xi \mathcal{J}_0$. In the non-covariant gauge we are choosing, this term is responsible for the Ward identity of three-point functions, but we do not expect it to contribute to higher-point functions.

4.2. Graviton

We consider the metric as a sum of the AdS metric and a graviton perturbation $g_{\mu\nu} = \frac{1}{x_0^2}(\delta_{\mu\nu} + h_{\mu\nu})$ and consider the action (see also [13])

$$I = \int d\alpha \, (R - 2\Lambda + \frac{1}{2} h_{\mu\nu} T^{\mu}_{\nu}) \ .$$

In the following we shall assume that there is no index raising for h, but for the energy-momentum tensor $T_{\mu\nu}$ the indices will be raised by $g_0^{\mu\nu} = x_0^2 \delta^{\mu\nu}$. We shall consider the gauge $h_{0\mu} = 0$. In this gauge, the linearised Einstein equations for h_{ij} become (see also [28])

$$\partial_0^2 (h_{ij} - \delta_{ij} h) - \frac{d-1}{x_0} \partial_0 (h_{ij} - \delta_{ij} h)$$
$$+ \partial_k^2 \tilde{h}_{ij} - \partial_l \partial_i \tilde{h}_{jl} - \partial_l \partial_j \tilde{h}_{il} + \delta_{ij} \partial_l \partial_m \tilde{h}_{lm} = -2T_{ij} \ , \tag{4.7}$$

where $h = \delta_{ij} h_{ij}$ and $\tilde{h}_{ij} = h_{ij} - \frac{1}{2} \delta_{ij} h$. There are also two constraints following from the 00 and $0i$ components of the Einstein equations,

$$\partial_0 (\partial_j h_{ij} - \partial_i h) = 2T_{0i} \ , \tag{4.8}$$

$$-\partial_i \partial_j h_{ij} + \partial_i^2 h - \frac{d-1}{x_0} \partial_0 h = 2T_{00} \ . \tag{4.9}$$

We decompose h_{ij} as

$$h_{ij} = \bar{h}^{\perp}_{ij} + \partial_i B^{\perp}_j + \partial_j B^{\perp}_i + \partial_i \partial_j \eta + \frac{1}{d-1} (\delta_{ij} - \frac{\partial_i \partial_j}{\partial^2}) h' \ ,$$

with $\partial^2 = \partial_i \partial_i$ and

$$B^{\perp}_i = \frac{1}{\partial^2} \partial_k h_{ik} - \frac{\partial_i}{(\partial^2)^2} \partial_k \partial_l h_{kl} \ , \qquad \eta = \frac{1}{(\partial^2)^2} \partial_k \partial_l h_{kl} \ , \qquad h' = h - \frac{1}{\partial^2} \partial_k \partial_l h_{kl} \ .$$

It is easy to check that

$$\delta_{ij} \bar{h}^{\perp}_{ij} = 0 \ , \qquad \partial_i \bar{h}^{\perp}_{ij} = 0 \ , \qquad \partial_i B^{\perp}_i = 0 \ .$$

Then the equations (4.7)–(4.8) become

$$(\partial_0^2 + \partial_k^2) \bar{h}^{\perp}_{ij} - \frac{d-1}{x_0} \partial_0 \bar{h}^{\perp}_{ij} = -2t_{ij} \ , \qquad t_{ij} \equiv P_{ijkl} T_{kl} \ , \tag{4.10}$$

$$\partial_0 B_i^{\perp} = \frac{1}{\partial^2}(2T_{0i} + \partial_0\partial_i h') \ , \qquad \partial_0 h' = -\frac{2}{\partial^2}\partial_j T_{0j} \ , \tag{4.11}$$

$$\partial_0 h = -\frac{2x_0}{d-1}T_{00} + \frac{x_0}{d-1}\partial^2 h' \ , \tag{4.12}$$

where P_{ijkl} is the transverse traceless projector in flat space.

Thus the equations for the transverse traceless part $\bar{h}_{ij}^{\perp} = P_{ijkl}h_{kl}$ reduce to that of a free massless scalar, so that we get ($d\beta = d^{d+1}y\sqrt{g_0(y)}$)

$$h_{ij}^{\perp}(x) = 2\int d\beta \ y_0^2 \ G(x,y) \ t_{ij}(y) \ . \tag{4.13}$$

Using the conservation of the energy-momentum tensor and the equations (4.11) and (4.12) we find that the quadratic (graviton propagator) part of the Einstein action takes the form

$$I = \frac{1}{2}\int d\alpha \ h_{ij}T_j^i = \int d\alpha d\beta \ (x_0 y_0)^2 \ t_{ij}(x) \ G(x,y) \ t_{ij}(y) - 2\int d\alpha \ x_0^2 \ T_{0i}\frac{1}{\partial^2}T_{0i}$$
$$- \frac{2}{d-1}\int d\alpha \ x_0^2 \ \partial_i T_{0i}\frac{1}{\partial^2}T_{00} - \frac{d-2}{d-1}\int d\alpha \ x_0^2 \ \partial_i T_{0i}(\frac{1}{\partial^2})^2\partial_i T_{0i} \ , \tag{4.14}$$

where we have ignored boundary terms resulting from partial integration.

5. Scalar four-point functions in AdS_5 supergravity / $\mathcal{N} = 4$ SYM theory

In this section we shall consider the four-point functions involving the scalar operators $\mathcal{O} = trF^2$ and $\mathcal{O}' = trFF^*$ in $\mathcal{N} = 4$ SYM theory in the large N and large 't Hooft coupling limit as predicted by the type IIB supergravity in $AdS_5 \times S^5$. $\mathcal{O}$ and $\mathcal{O}'$ correspond to the massless fields ϕ and C in AdS_5 coming from the dilaton and axion of type IIB supergravity on $AdS_5 \times S^5$. The four-point functions of interest are $< \mathcal{O}(x_1)\mathcal{O}(x_2)\mathcal{O}(x_3)\mathcal{O}(x_4) >$, $< \mathcal{O}'(x_1)\mathcal{O}'(x_2)\mathcal{O}(x_3)\mathcal{O}(x_4) >$ and $< \mathcal{O}'(x_1)\mathcal{O}'(x_2)\mathcal{O}'(x_3)\mathcal{O}'(x_4) >$. They correspond in AdS_5 to the tree-level scattering amplitudes of four dilatons, two dilatons and two axions, and four axions respectively. Thus we need to look for vertices in the Lagrangian of IIB supergravity on $AdS_5 \times S^5$ involving two dilatons (axions) and one other field or quartic vertices involving only dilatons and axions. For this purpose it is sufficient to concentrate on the graviton-dilaton-axion sector of the supergravity[8]

$$\hat{I} = \frac{1}{2\kappa_{10}^2}\int_{AdS_5} d^5x \int_{S^5} d^5y \ \sqrt{-\hat{g}} \ \left[\hat{R} - \frac{1}{2}(\partial\hat{\phi})^2 - \frac{1}{2}e^{2\hat{\phi}}(\partial\hat{C})^2\right] . \tag{5.1}$$

[8] Our notation will be as follows. All fields in $D = 10$ will be written with hats on them and their indices. $\mu, \nu, ...$ will refer to the indices of AdS_5, while $\alpha, \beta, ...$ will refer to those of S^5.

Since the fields ϕ and C are associated with the zeroth spherical harmonics of $\hat{\phi}$ and $\hat{C}$ on S^5, it is clear that the desired vertices cannot involve fields from higher spherical harmonics. Thus the fields in (5.1) can be assumed to have only x dependence, i.e.

$$\hat{\phi}(x,y)=\phi(x), \qquad \hat{C}(x,y)=C(x), \qquad \hat{h}_{\hat{\mu}\hat{\nu}}(x,y)=\{h_{\mu\nu}(x),h_{\mu\alpha}(x),h_{\alpha\beta}(x)\}\ .$$

After dimensional reduction to AdS_5 and a Weyl scaling[9] the relevant part of (5.1) becomes

$$I=\frac{1}{2\kappa^2}\int_{AdS_5} d^5x\sqrt{-g'}\ \left[R-\frac{1}{2}(\partial\phi)^2-\frac{1}{2}e^{2\phi}(\partial C)^2\right]\ , \tag{5.2}$$

where $R=R(h')$ is the 5d Ricci scalar. The $\phi-C$ sector of the IIB supergravity on $AdS_5\times S^5$ is thus very simple. The only cubic vertex involving two ϕ is $\partial\phi\partial\phi\ h'$. In particular, there is no $h^\alpha_\alpha\phi\phi$ vertex, where h^α_α is the trace of the internal (S^5) part of the metric (massive fixed scalar) which in the $\mathcal{N}=4$ SYM theory corresponds to the operator of the structure $\mathcal{O}_8=trF^4-\frac{1}{4}tr(F^2)^2$. Since in the free Maxwell theory there is a non-vanishing three-point function $<\mathcal{O}\mathcal{O}\mathcal{O}_8>$ [29], this three-point function must have non-trivial dependence on the 't Hooft coupling (cf. [17]).

Following the same reasoning, we can see that the diagrams contributing to the four-graviton scattering may come only from the R term in (5.2). In particular, there is no cubic vertex involving two gravitons and other fields. This is consistent with the expectation [30] that the OPE of the energy momentum tensor in $\mathcal{N}=4$ SYM theory closes on itself.[10]



Figure 2: Scattering diagrams for $\phi\phi\phi\phi$ and $CCCC$. Only s-channel diagrams are displayed here.

[9] The resulting graviton $h'_{\mu\nu}$ differs from $h_{\mu\nu}$ by a Weyl scaling, i.e. $h'_{\mu\nu}=h_{\mu\nu}+\frac{1}{3}g_{0\mu\nu}h^\alpha_\alpha$, where $g_{0\mu\nu}$ is the background metric in AdS_5.

[10] This is not so in most of four-dimensional theories. For a discussion of the case of the $\mathcal{N}=1$ supersymmetric theories see [31].



Figure 3: Scattering diagrams for $\phi\phi CC$.

Starting with (5.2), we can write down the scattering diagrams in AdS_5 contributing to the four-point functions under consideration (see Fig.2 and Fig.3). Let us first look at the diagrams which do not involve graviton exchange. The scattering amplitude with the dilaton or axion exchange, e.g., $\phi(1)C(2)\phi(3)C(4)$ in Fig.3, can be written as

$$A_1 = \int \frac{du_0 d^d u}{u_0^{d+1}} \frac{dv_0 d^d v}{v_0^{d+1}} (u_0 v_0)^2 \partial_\mu \mathcal{K}_d(u, x_1) \partial_\mu \mathcal{K}_d(u, x_2) G(u, v) \partial_\nu \mathcal{K}_d(v, x_3) \partial_\nu \mathcal{K}_d(v, x_4) \ . \tag{5.3}$$

The contact $\phi C\phi C$ interaction gives

$$A_2 = \int \frac{du_0 d^d u}{u_0^{d+1}} \, u_0^2 \, \mathcal{K}_d(u, x_1) \partial_\nu \mathcal{K}_d(u, x_2) \mathcal{K}_d(u, x_3) \partial_\nu \mathcal{K}_d(u, x_4) \ . \tag{5.4}$$

Here $\mathcal{K}_d$ ($d = 4$) and $G(u, v)$ (see (3.2) and (3.3)) are the boundary and bulk propagators for the massless scalars. Since these propagators satisfy

$$\frac{1}{\sqrt{g_0}} \partial_\mu (\sqrt{g_0} g_0^{\mu\nu} \partial_\nu) \mathcal{K}_d = 0 \ , \qquad \frac{1}{\sqrt{g_0}} \partial_\mu (\sqrt{g_0} g_0^{\mu\nu} \partial_\nu) G(u, v) = -\frac{1}{\sqrt{g_0}} \delta(u - v) \ ,$$

we find by a partial integration[11] that $A_1(1, 2, 3, 4)$ reduces to $-\frac{1}{2} A_2(1, 3, 2, 4)$.

Direct evaluation of A_2 is quite tedious. It is easy to show, however, that the symmetric in $x_1, ..., x_4$ part of (5.4) $(A_2)_{sym}$ (and thus $(A_1)_{sym}$) vanishes. One way to see that is to make a field redefinition, $\phi = -\log(1 + \psi)$. Then the free action for ϕ becomes

$$I = \int d^{d+1} x \sqrt{g_0} \ (1 + \psi)^{-2} (\partial \psi)^2 \ . \tag{5.5}$$

The new action generates cubic and quartic dilaton vertices and it can be seen that the resulting four-dilaton scattering amplitude is proportional to $(A_2)_{sym}$. Since the field

[11] Boundary terms resulting from partial integration give only unimportant contact terms (terms containing Dirac delta-function and its derivatives).

redefinition changes the correlators only by unimportant contact terms, we conclude that $(A_2)_{sym}$ should contain only contact terms, i.e. should vanish for separated points.[12]

The fact that $(A_2)_{sym} = -2(A_1)_{sym} = 0$ implies, in particular, that as in the case of the $\phi\phi\phi\phi$ amplitude, the only non-trivial contribution to the $CCCC$ amplitude comes from the sum of the s, t, u graviton exchanges, so that the $\phi\phi\phi\phi$ and $CCCC$ amplitudes are actually equal.[13] The structure of the $\phi C\phi C$ amplitude is more complicated as in addition to the graviton exchange it contains also the sum of the axion exchange and the contact contributions given by a combination of A_2 amplitudes with different orders of end-points (see also [32]).

The formal integral expression for the graviton contribution to the four-point functions can be written down by substituting into (4.14) the appropriate expression for the energy-momentum tensor corresponding to the massless scalar fields,

$$T_{\mu\nu} = \partial_\mu \mathcal{K}_d(u, x_1)\, \partial_\nu \mathcal{K}_d(u, x_2) - \frac{1}{2}\delta_{\mu\nu}\, \partial_\lambda \mathcal{K}_d(u, x_1)\, \partial_\lambda \mathcal{K}_d(u, x_2)\ .$$

Here $\mathcal{K}_d$ is again the boundary propagator (with $d = 4$). The resulting expression is quite long and will not be explicitly presented here. We leave its detailed analysis for future work.

To conclude, let us make some speculative remarks on consequences of possible crossing symmetry of CFT four-point functions for the structure of scattering amplitudes in AdS. From correspondence between the scattering amplitudes in AdS and correlators in CFT given by CPWE's, we would expect that the amplitudes in Fig.2 and Fig.3 should possess $s - t - u$ symmetry. That would imply, for example, that the s, t and u channel graviton exchanges in the $\phi\phi\phi\phi$ amplitude are all equal. Then in the $CCCC$ amplitude (which does not contain contact contribution) the s, t, u dilaton exchages would also need to be equal. Since their sum $(A_1)_{sym}$ vanishes, each of them should also be vanishing, at least

[12] In the original version of this paper we were assuming that this argument demonstrates the triviality of A_2, but it actually applies only to $(A_2)_{sym}$ as the the four legs here correspond to the *same* field (while $\psi^2(\partial\psi)^2$ reduces after integration by parts to $-\frac{1}{3}\psi^3\partial^2\psi$ that vanishes on shell, there is no similar rearrangement for $\phi^2(\partial C)^2$). We noticed this mistake after reading the recent paper [32] with conclusions of which we now agree.

[13] This may be viewed as a consequence of the $SL(2, R)$ symmetry of the action (5.2).

up to some singular contributions (cf. [32]) that would need to be subtracted as part of a definition of the CFT/AdS correspondence.[14] To be able to draw any conclusions from the assumption of crossing symmetry in the case of the $\phi C\phi C$ amplitude (Fig.3) one needs first to interpret the contact bulk diagram contributing to it. For example, we may replace it by a massless scalar exchange in the t-channel using that, as shown above, $A_2(1,2,3,4) = -2A_2(1,3,2,4)$.[15] Given that in $\phi C\phi C$ case the graviton exchange contributes only to the t-channel, one would be led to the conclusion that the scalar and graviton exchange amplitudes should be proportional to each other (up to possible subtractions mentioned above). That would, in turn, imply that all these massless scalar four-point amplitudes should be vanishing. Given these somewhat surprising conclusions, it remains to be seen if the crossing symmetry can actually be realised in a four-dimensional CFT.

Appendix A. Scalar propagator in AdS_{d+1}

As in [6] we consider the Anti de Sitter space of dimension $D = d+1$ with Euclidean signature and the (half-space) metric

$$ds^2 = g_{0\mu\nu}dx^\mu dx^\nu = \frac{1}{x_0^2}(dx_0^2 + dx_i^2) \ . \tag{A.1}$$

The AdS_{d+1} bulk indices will be denoted by $\mu, \nu, ...$ and will take values $0, 1, ..., d$. We shall use the notation $x = (x_0, \vec{x})$, $\vec{x} = (x_i)$, $\ i = 1, ..., d$.

Considering the Euclidean action for a massive scalar

$$I = \frac{1}{2}\int d^{d+1}x\sqrt{g_0}\,[(\partial_\mu\phi)^2 + m^2\phi^2] \ ,$$

[14] Note that in Euclidean space the massless scalar exchange diagrams are generated by a contact-type (δ-function) vertex $< \mathcal{O}(x)\mathcal{O}'(y)\mathcal{O}'(z) > \sim [\delta(x-y)+\delta(x-z)]\frac{1}{|y-z|^8} - \delta(y-z)\frac{1}{|x-z|^8}$ represented in the bulk by the $\phi\partial C\partial C$ interaction (its non-contact part vanishes since upon integration by parts it reduces to $\frac{1}{2}C^2\partial^2\phi - \phi C\partial^2 C$). However, the corresponding Minkowskian (Wightman) three-point function is zero and thus should not contribute to the CPWE of the four-point Wightman function in CFT.

[15] The massless scalar here may be identified with dilaton, as there is a non-vanishing contact-type Euclidean three-point function in SYM theory: $< \mathcal{O}(x)\mathcal{O}(y)\mathcal{O}(z) > \sim \delta(x-y)\frac{1}{|y-z|^8} + \delta(x-z)\frac{1}{|y-x|^8} + \delta(y-z)\frac{1}{|x-z|^8}$. This is also suggested by the field redefinition argument in (5.5).

we are to solve

$$(D^2 - m^2)G(x,y) = -\frac{1}{\sqrt{g_0}}\delta(x-y) \ .$$

More explicitly, in the coordinate system (A.1), this equation becomes:

$$x_0^n \partial_\mu [x_0^{-n+2}\partial_\mu G(x,y)] - m^2 G(x,y) = -\delta(\vec{x}-\vec{y})\delta(x_0-y_0)y_0^n \ .$$

Let $G(x,y) = x_0^{d/2}H(x,y)$. Then

$$(\Delta_\nu + \partial_i^2)H(x,y) = -y_0^{\frac{d-2}{2}}\delta(\vec{x}-\vec{y})\delta(x_0-y_0) \ , \tag{A.2}$$

where $\nu^2 = m^2 + \frac{1}{4}d^2$ and the operator Δ_ν defined by

$$\Delta_\nu = \partial_0^2 + \frac{1}{x_0}\partial_0 - \frac{\nu^2}{x_0^2} \ ,$$

has Bessel functions as its eigenvalues

$$\Delta_\nu J_\nu(wx_0) = -w^2 J_\nu(wx_0) \ .$$

The delta-functions can be written in terms of orthonormal functions:

$$\delta(x_0-y_0) = y_0\int_0^\infty dw\, wJ_\nu(wx_0)J_\nu(wy_0), \qquad \int \frac{d^dk}{(2\pi)^d}e^{i\vec{k}\cdot(\vec{x}-\vec{y})} = \delta(\vec{x}-\vec{y}) \ .$$

We expand the Green function in a similar fashion:

$$H(x,y) = \int \frac{d^dk}{(2\pi)^d}e^{i\vec{k}\cdot\vec{x}}\int_0^\infty dw\, wJ_\nu(wx_0)\tilde{H}(w,k;y) \ .$$

Then the substitution into (A.2) leads to

$$\tilde{H}(w,k;y) = y_0^{d/2}\frac{J_\nu(wy_0)}{w^2+k^2}e^{-i\vec{k}\cdot\vec{y}} \ .$$

The scalar Green function in the Anti-de Sitter space (A.1) is thus given by

$$\begin{aligned} G(x,y) &= (x_0y_0)^{d/2}\int\frac{d^dk}{(2\pi)^d}\int_0^\infty dw\ w\frac{1}{w^2+k^2}e^{i\vec{k}\cdot(\vec{x}-\vec{y})}J_\nu(wx_0)J_\nu(wy_0) \\ &= (x_0y_0)^{d/2}\int\frac{d^dk}{(2\pi)^d}\,e^{i\vec{k}\cdot(\vec{x}-\vec{y})}I_\nu(kx_0^<)K_\nu(kx_0^>) \ , \end{aligned} \tag{A.3}$$

where $x_0^<(x_0^>)$ is the smaller (larger) number among x_0 and y_0.

Appendix B. Scalar contribution to four-point functions

The contribution of a scalar operator Φ of dimension λ to the CPWE of a four-point function of scalar operators $< 0|\Phi_1(x_1)\Phi_2(x_2)\Phi_3(x_3)\Phi_4(x_4)|0 >$ has been worked out long time ago in [19]. The same expression should also correspond to the scattering amplitude in AdS (3.1). Here we shall quote the result of [19]. We will use the following definitions: $\Sigma_{ij} = \lambda_i + \lambda_j$, $\Delta_{ij} = \lambda_i - \lambda_j$, $x_{ij} = |x_i - x_j|$, and

$$\rho = \frac{x_{14}^2 x_{23}^2}{x_{12}^2 x_{34}^2}, \qquad \eta = \frac{x_{13}^2 x_{24}^2}{x_{12}^2 x_{34}^2} \ ,$$

where $\lambda_i,\ i = 1, 2, 3, 4$ are the conformal dimensions of Φ_i. Then

$$< 0|\Phi_1(x_1)\Phi_2(x_2)\Phi_3(x_3)\Phi_4(x_4)|0 >= x_{12}^{\Sigma_{12}-\Delta_{12}} x_{13}^{-\Delta_{12}-\Delta_{34}} x_{14}^{\Delta_{34}-\Delta_{12}} x_{34}^{\Delta_{12}-\Sigma_{34}} f_0(\rho,\eta) \ ,$$

with

$$f_0(\rho,\eta) = c_1 \eta^{-\frac{1}{2}(\lambda-\Delta_{12})} \Big[F_4\big(\frac{1}{2}(\lambda+\Delta_{34}), \frac{1}{2}(\lambda-\Delta_{12}), \lambda+1-\frac{1}{2}d, \frac{1}{2}(\Delta_{34}-\Delta_{12})+1; \frac{1}{\eta}, \frac{\rho}{\eta}\big)$$

$$+ c_2 (\frac{\rho}{\eta})^{-\frac{1}{2}(\lambda-\Delta_{12})} F_4\big(\frac{1}{2}(\lambda-\Delta_{34}), \frac{1}{2}(\lambda-\Delta_{12}), \lambda+1-\frac{1}{2}d, 1-\frac{1}{2}(\Delta_{34}+\Delta_{12}); \frac{1}{\rho}, \frac{\eta}{\rho}\big) \Big]$$

When $\Delta_{12} = \Delta_{34} = 0$, this simplifies into

$$f_0(\rho,\eta) = c_1' \eta^{-\frac{1}{2}\lambda} \Big[F_4\big(\frac{1}{2}\lambda, \frac{1}{2}\lambda, \lambda+1-\frac{1}{2}d, 1; \frac{1}{\eta}, \frac{\rho}{\eta}\big)$$

$$+ c_2' (\frac{\rho}{\eta})^{-\frac{1}{2}\lambda} F_4\big(\frac{1}{2}\lambda, \frac{1}{2}\lambda, \lambda+1-\frac{1}{2}d, 1; \frac{1}{\rho}, \frac{\eta}{\rho}\big) \Big] \ .$$

In above c_1, c_2, c_1', c_2' are some numerical constants and F_4 is a double hypergeometric function.

Acknowledgments

We acknowledge the support of PPARC and of the European Commission TMR programme grant ERBFMRX-CT96-0045. We are grateful to S. Ferrara, D. Freedman, I. Klebanov, R. Metsaev and A. Ritz for stimulating discussions.

References

[1] G. 't Hooft, "A Planar diagram theory for strong interactions", Nucl. Phys. B72 (1974) 461.

[2] A.M. Polyakov, "String theory and quark confinement", hep-th/9711002; "A few projects in string theory", Les Houches Summer School, 1992:783, hep-th/9304146.

[3] J. Maldacena, "The large N limit of superconformal field theories and supergravity", hep-th/9711200.

[4] I.R. Klebanov, "World-volume approach to absorption by non-dilatonic branes", Nucl. Phys. B496 (1997) 231, hep-th/9702076; S.S. Gubser, I.R. Klebanov and A.A. Tseytlin, "String theory and classical absorption by three-branes", Nucl. Phys. B499 (1997) 41, hep-th/9703040; S.S. Gubser and I.R. Klebanov, "Absorption by branes and Schwinger terms in the world-volume theory", Phys. Lett. B413 (1997) 41, hep-th/9708005.

[5] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, "Gauge theory correlators from non-critical string theory", hep-th/9802109.

[6] E. Witten, "Anti de Sitter space and holography", hep-th/9802150.

[7] G. Horowitz and H. Ooguri, "Spectrum of Large N Gauge Theory from Supergravity", Phys. Rev. Lett. 80 (1998) 4116, hep-th/9802116.

[8] S. Ferrara, C. Fronsdal and A. Zaffaroni, "On N=8 Supergravity on AdS_5 and $N = 4$ Superconformal Yang-Mills theory", hep-th/9802203.

[9] I.Ya. Aref'eva and I.V. Volovich, "On large N conformal theories, field theories in Anti de Sitter space and singletons", hep-th/9803028; "On the Breaking of Conformal Symmetry in the AdS/CFT Correspondence", hep-th/9804182.

[10] W. Mück and K.S. Viswanathan, "Conformal field theory correlators from classical scalar field theory on AdS_{d+1}", hep-th/9804035.

[11] M. Henningson and K. Sfetsos, "Spinors and the AdS/CFT correspondence", hep-th/9803251.

[12] D.Z. Freedman, S.D. Mathur, A. Matusis and L. Rastelli, "Correlation functions in the $\mathrm{CFT}_d/\mathrm{AdS}_{d+1}$ correspondence", hep-th/9804058.

[13] H. Liu and A. A. Tseytlin, "D=4 Super-Yang-Mills, D=5 Gauged Supergravity, and D=4 Conformal Supergravity," hep-th/9804083.

[14] G. Chalmers, H. Nastase, K. Schalm and R. Siebelink, "R-Current Correlators in $\mathcal{N} = 4$ SYM from AdS," hep-th/9805015.

[15] W. Mück and K. S. Viswanathan, "Conformal Field Theory Correlators from Classical Field Theory on Anti-de Sitter Space II. Vector and Spinor Fields," hep-th/9805145.

[16] A. M. Ghezelbash, K. Kaviani, S. Parvizi and A. H. Fatollahi, "Interacting Spinors-Scalars and the AdS/CFT Correspondence," hep-th/9805162.

[17] S. Lee, S. Minwalla, M. Rangamani and N. Seiberg, "Three-Point Functions of Chiral Operators in D=4, $\mathcal{N} = 4$ SYM at Large N", hep-th/9806074.

[18] D.Z. Freedman, S.D. Mathur, A. Matusis and L. Rastelli, to appear, talk by D.Z. Freedman at Strings 98, Santa Barbara.

[19] S. Ferrara, R. Gatto, A.F. Grillo and G. Parisi, Nucl. Phys. B49 (1972) 77; Nuovo Cimento 26 (1975) 226.

[20] S. Ferrara, R. Gatto, A.F. Grillo and G. Parisi, Nuovo Cim. Lett. 4 (1972) 115; Nuovo Cim. Lett. 5 (1972) 147; A. M. Polyakov, JETP 39 (1974) 10; M. Ya. Palchik, Phys. Lett. 66B (1977) 259; G. Mack Commun. Math. Phys. 53 (1977) 155.

[21] E.S. Fradkin and M.Ya. Palchik, "Conformal Quantum Field Theory in D dimensions" (Kluwer, Dordrecht, 1996).

[22] E.S. Fradkin and M.Ya. Palchik, Phys. Rept. 44C (1978) 249.

[23] I.T. Todorov, M.C. Mintchev and V.B. Petkova "Conformal Invariance in Quantum Field Theory (Scuola Normale Superiore, Pisa, 1978).

[24] B. Allen and T. Jacobson, "Vector two-point functions in maximally symmetric spaces", Commun. Math. Phys. 103 (1986) 669.

[25] B. Allen, "Graviton propagator in de Sitter space", Phys. Rev. D 34 (1986) 3670; B. Allen and M. Turyn, "An evaluation of the graviton propagator in de Sitter space" Nucl. Phys. B292 (1987) 813.

[26] T. Banks and M.B. Green, "Non-perturbative Effects in $AdS_5 \times S_5$ String Theory and d=4 SUSY Yang-Mills", hep-th/9806004; M. Bianchi, M.B. Green, S. Kovacs and G. Rossi, "Instantons in supersymmetric Yang-Mills and D-instantons in IIB superstring theory", hep-th/9807033.

[27] S. Ferrara and A. Zaffaroni, "Bulk gauge fields in AdS supergravity and supersingletons", hep-th/9807090.

[28] G.E. Arutyunov and S. Frolov, "On the origin of supergravity boundary terms in the AdS/CFT correspondence", hep-th/9806216.

[29] S.S. Gubser, A. Hashimoto, I.R. Klebanov and M. Krasnitz, "Scalar absorption and the breaking of the world volume conformal invariance", hep-th/9803023; S. S. Gubser and A. Hashimoto, "Exact absorption probabilities for the D3-brane", hep-th/9805140.

[30] P.S. Howe and P.C. West, "Operator product expansions in four-dimensional superconformal field theories", Phys. Lett. B389 (1996) 273, hep-th/9607060.

[31] D. Anselmi, M. Grisaru and A. Johansen, "A Critical Behaviour of Anomalous Currents, Electric-Magnetic Universality and CFT_4", hep-th/9601023, Nucl.Phys. B491 (1997) 221.

[32] D.Z. Freedman, S.D. Mathur, A. Matusis and L. Rastelli, "Comments on 4-point functions in the CFT/AdS correspondence", hep-th/9808006.